ERRATUM Open Access



Erratum to: Characterization of a Plasmodium berghei sexual stage antigen PbPH as a new candidate for malaria transmission-blocking vaccine

Xu Kou^{1,2}, Wenqi Zheng¹, Feng Du¹, Fei Liu³, Meilian Wang¹, Qi Fan⁴, Liwang Cui⁵, Enjie Luo^{1*} and Yaming Cao^{3*}

Erratum

After the publication of our paper [1], we realized that the positions of the primers 2, 5 and 6 in Fig. 4a were incorrect due to shifting. The corrected Fig. 4 is included below. In addition, "Mosquitoes Infected/Dissected" in Table 1 should read "Mosquitoes Dissected/Infected". We would like to apologise for these errors and for any inconvenience this may have caused.

Author details

¹Department of Pathogen Biology, College of Basic Medical Sciences, China Medical University, Shenyang, Liaoning 110001, China. ²College of Animal Husbandry and Veterinary, Liaoning Medical University, Jinzhou, Liaoning 121001, China. ³Department of Immunology, College of Basic Medical Sciences, China Medical University, Shenyang, Liaoning 110001, China. ⁴Dalian Institute of Biotechnology, Dalian, Liaoning, China. ⁵Department of Entomology, The Pennsylvania State University, University Park, PA 16802, ISA

Received: 6 February 2017 Accepted: 6 February 2017 Published online: 16 February 2017

Reference

 Kou X, Zheng W, Du F, Liu F, Wang M, Fan Q, et al. Characterization of a Plasmodium berghei sexual stage antigen PbPH as a new candidate for malaria transmission-blocking vaccine. Parasit Vectors. 2016;9:190. doi:10.1186/s13071-016-1459-8.

^{*} Correspondence: enjieluo@hotmail.com; ymcao@mail.cmu.edu.cn

1 Department of Pathogen Biology, College of Basic Medical Sciences, China Medical University, Shenyang, Liaoning 110001, China

3 Department of Immunology, College of Basic Medical Sciences, China Medical University, Shenyang, Liaoning 110001, China






Fig. 4 Knockout of *pbph* gene from *P. berghei* parasite. **a** A schematic shows the wild type *pbph* locus, transfection construct, and the recombined locus. Through a double-crossover strategy the *pbph* locus was replaced with the *dhfr*-expressing cassette. Primers 1–6 used to detect gene knockout are marked. **b** PCR detection of wild-type and *Δpbph* parasite genomes. Lane 1: primers 1 + 2 (910 bp); Lane 2: primers 1 + 3 (990 bp); Lane 3: primers 5 + 6 (1,040 bp)